

## Distribution and characteristics of *Vibrio vulnificus* in Incheon Coastal Area.

Y.H.Kim, J.M Koh, J.M. Lee, Y.W. Gong, B.Y. Oh, M.Y. LEE, H.Y.Kim, K.W Hwang, Y.J Koh, J.H.Kim, S.Jegal

Incheon Institute of Health & Environment, Incheon, Korea

*Vibrio vulnificus*, a gram-negative, and estuarine bacterium commonly found in coastal water and in association with seawater, shellfish and fish. This bacterium is known to cause septicemia and severe wound infections in patients with chronic liver disease or immuno-compromised condition. In this study, we investigated the distribution and characteristics of *Vibrio vulnificus* in Incheon Coastal Area.

In this survey, total 6,571 samples were obtained from different sites of the Incheon coastal area during the periods from 2003 to 2005. 1,272 strains were isolated from 6,571 samples by PCR using targeted the cytotoxin-hemolysin gene. The isolation rates of *Vibrio vulnificus* from sediment, seawater, raw seafoods and aquarium water were 40.6%, 25.9%, 5.7%, and 29.4%, respectively. The highest isolation rate was 31.8% in September 2005 and in spite of low temperature the isolation rate was 13.1% in November 2005.

These results show that the prevalence of *Vibrio vulnificus* was highest in the sediment among environment samples. Thus, we should recognize sediment plays role in as potential reservoir of *Vibrio vulnificus*



# Distribution and characteristics of *Vibrio vulnificus* in Incheon Coastal Area.

Y.H.Kim, J.M Koh, J.M. Lee, Y.W. Gong, B.Y. Oh, M.Y. LEE,  
H.Y.Kim, K.W Hwang, Y.J Koh, J.H.Kim, S.Jegal  
Incheon Institute of Health & Environment, Incheon, Republic of Korea

## Abstract

**Background:** *Vibrio vulnificus*, a gram-negative, and estuarine bacterium commonly found in coastal water and in association with estuarine water, shellfish and fish. This bacterium is known to cause septicemia and severe wound infections in patients with chronic liver disease or immuno-compromised condition. In this study, we investigated the distribution and characteristics of *Vibrio vulnificus* in Incheon Coastal Area. **Method:** In this survey, total 4,302 samples were obtained from different sites of the Incheon coastal area during the periods from 2003 to 2005. 310 strains were isolated from 4,302 samples by biochemical test. The isolation rates of *Vibrio vulnificus* from sediment, estuarine water and Fishes & Shellfishes were 23.2%, 12.4% and 4.7%, respectively. The highest isolation rate was 31.8% in September 2005 and in spite of low temperature the isolation rate was 13.1% in November 2005. **Conclusion:** These results show that the prevalence of *Vibrio vulnificus* was highest in the sediment among environment samples. Thus, we should recognize sediment plays role in as potential reservoir of *Vibrio vulnificus*.

## Introduction

*Vibrio vulnificus*, a gram-negative, and estuarine bacterium commonly found in coastal water and in association with estuarine water, shellfish and fish. This bacterium is known to cause septicemia and severe wound infections in patients with chronic liver disease or immuno-compromised condition. Primary septicemia usually occurs through ingestion of raw shellfish, especially oysters, by persons who are immunocompromised. Wound infections however, have been known to occur in otherwise healthy individuals who come in contact with this bacterium via contamination of a previously inflicted wound or one incurred in an estuarine environment. One of the main virulence factors associated with infection with *V. vulnificus* is an antiphagocytic, polysaccharide capsule, and encapsulated cells are highly virulent, with a 50% lethal dose of less than 10 CFU. The *V. vulnificus* endotoxin is also important in the virulence of this pathogen and the hypotension produced by this factor is eliminated when nitric oxide synthase is inhibited. In this report, we investigated the distribution and characteristics of *Vibrio vulnificus* in Incheon Coastal Area, which were isolated from total 4,302 samples (sediment, estuarine water, fishes & shellfishes) in 2004-2006.

## Methods



Fig.1. Schematic view of this study

Fig. 2. Sampling sites for the isolation of *V. vulnificus*





Fig.3. Isolation rates of *Vibrio vulnificus* during three years




Fig.4. Isolation rates of *Vibrio vulnificus* from marine specimens




Fig.5. Correlation between isolation rates of *Vibrio vulnificus* and mean temperatures of estuarine water




Fig.6. Correlation between isolation rates of *Vibrio vulnificus* and mean salinity of estuarine water

## Results

- We investigated, total 4,302 samples were obtained from different sites of the Incheon coastal area during the periods from 2004 to 2006. 310 strains were isolated from 4,302.
- The isolation rates of *Vibrio vulnificus* from sediment, estuarine water and fishes & shellfishes were 23.2%, 12.4% and 4.7% respectively.
- The isolation rates of *Vibrio vulnificus* were positively correlated with water temperature and negatively correlated with salinity.

Table 1. Isolation rates of *Vibrio vulnificus* in Incheon coastal area

Specimen	No. of sample (N)	No. of isolated (n)	(%)
Fishes & Shellfishes	3,258	153	(4.7)
Estuarine water	792	98	(12.4)
Sediment	254	59	(23.2)
Total	4,302	310	(7.2)

## Conclusion

These results suggest that the prevalence of *Vibrio vulnificus* was highest in the sediment among environment samples. Thus, we should recognize sediment plays role in as potential reservoir of *Vibrio vulnificus*.



Incheon Institute of Health & Environment

## References

1. Ecology of *Vibrio vulnificus* in Estuarine waters of Eastern North Carolina. CS Pfeiffer, MF Hite, JD Oliver. *Applied and Environmental Microbiology*. 2003; 3526-3531.
2. Influence of Water Temperature and Salinity on *Vibrio vulnificus* in northern Gulf and Atlantic Coast Oyster. M.L.MOTES, A.DuPAULA, D.W.COOK, J.E.VEAZEY, J.C.HUNSLUCKER. *Applied and Environmental Microbiology*. 1996; 1459-1465.
3. Environmental and Antimicrobial Characteristics of *Vibrio* spp. Isolated from Fish, Shellfish, Seawater and Brackish water samples in Gyeongju, Eastern Coast, Korea. *Korean Journal of Environmental Health*. 2003; 29: 94-102.
4. Kunmerer, K. Significance of antibiotics in the environment. *Journal of Antimicrobial Chemistry*. 52: 6-10, 2003.
5. Effects of Temperature and Salinity on *Vibrio vulnificus* Population Dynamics as Assessed by Qualitative PCR. Mek A. Hamed, Martin P. Pitz, Elin Lim. *Applied and Environmental Microbiology*. 1996; 1459-1465.
6. In vitro susceptibility of *Vibrio* spp. isolated from the environment. S.Zanten, T.Spau, A.Derks, L.Romano, L.A.Sechi, O.Fadda. *International Journal of Antimicrobial Agents*. 17: 2001; 407-409.
7. Detection and differentiation of *Vibrio vulnificus* in seawater and plankton of a coastal zone of the Mediterranean Sea. Teresa L.Mauger, Maria Carbone, Maria T. Fera, Concetta Gugliandolo. *International Journal of Antimicrobial Agents*. 2005; 1-7.